

JOURNAL OF APPROXIMATION THEORY 26, 101–107 (1979)

On Simultaneous Best Approximations in $C^1[a, b]$

M. R. SUBRAHMANYA

*Fachbereich Mathematik, Universität Frankfurt, 6 Frankfurt, West Germany**Communicated by P. L. Butzer*

Received April 18, 1977

INTRODUCTION

Brosowski [2] has proved the following:

THEOREM. *Let $V_1 \subset V_2$ be alternation systems of $C[a, b]$, $v_1 \in V_1$ and $v_2 \in V_2$. In order that there exists an element $f \in C[a, b]$ such that $v_i \in P_{V_i}(f)$, $i = 1, 2$, it is necessary and sufficient that $v_1 - v_2$ is either zero or changes sign at least $d(v_1)$ points of $[a, b]$. If V_1 and V_2 are contained in $C^2[a, b]$, then we can choose f to be a polynomial.*

For the case of polynomials, Rivlin [7] posed the following:

PROBLEM. *Characterize those n -tuples of algebraic polynomials $\{p_0, p_1, \dots, p_{n-1}\}$ with degree of $p_i = i$, $i = 0, 1, \dots, n-1$, for which there exists an f in $C[a, b]$ such that the polynomial of best approximation of degree i to f is p_i , $i = 0, 1, \dots, n-1$.*

He has shown that for this to be true it is necessary for each pair i, j with $0 \leq i < j \leq n-1$ that the polynomial $p_i - p_j$ is either zero or changes sign at least $i+1$ points of $[a, b]$. Deutsch, Morris, and Singer [5] have shown that in the case $n = 2$, the condition is also sufficient. Sprecher [8] has extended this result to the case of two polynomials of arbitrary degrees and in [9] has given a solution to the above problem for $n = 3$. Hegering [6] and Subrahmanya [10] have considered general Chebyshev systems of $C[a, b]$. Subrahmanya [11] has given a solution to the Rivlin problem for a general n . Brosowski and Subrahmanya [3] have considered $C[T]$, where T is a compact Hausdorff space and have characterized an infinite set of elements for which there exists an $f \in C[T]$ with this set of elements as best approximations from arbitrary subsets which they assume satisfy only in the necessity part. Hegering [6] has also proved the following:

Let $V_1 \subset V_2$ be Haar subspaces of $C[a, b]$ and $v_i \in V_i$, $i = 1, 2$. Let

$v_1 \in C^h[a, b]$ and $v_2 \in C^l[a, b]$, $h, l \in \mathbb{N}$ ($h, l = \infty$ allowed). Under the necessity condition of Rivlin there exists an $f \in C^{\min(h, l)}[a, b]$ such that $v_i \in P_{V_i}(f)$, $i = 1, 2$.

In this paper we generalize this result to alternation systems. Our proof is simple and straight forward.

We now give a brief review of some notation. Let $C[a, b]$ denote the set of all continuous real-valued functions defined on $[a, b]$. $C[a, b]$ is equipped with the norm given by

$$\|f\| := \sup_{t \in [a, b]} |f(t)|.$$

Let $V \subset C[a, b]$ and $f \in C[a, b]$. An element $v_0 \in V$ is said to be a best approximation to f in V , if

$$\|f - v_0\| = \inf_{v \in V} \|f - v\| =: E_V(f).$$

We denote by $P_V(f)$ the set of all best approximations to f from V , that is,

$$P_V(f) = \{v \in V \mid \|f - v\| = E_V(f)\}$$

A nonempty subset V of $C[a, b]$ is said to be an alternation system if there exists a mapping

$$d: V \rightarrow \mathbb{R}$$

such that $d(v) \geq 1$ for each $v \in V$ and if the following condition is fulfilled:

An element v in V is a best approximation of an f in $C[a, b]$ if and only if there exist points $t_0, t_1, \dots, t_{d(v)}$ with $a \leq t_0 < t_1 < \dots < t_{d(v)} \leq b$, and an $\eta \in \{-1, +1\}$ such that

$$\|f - v\| = \eta(-1)^k (f(t_k) - v(t_k)),$$

$k = 0, 1, \dots, d(v)$.

THE MAIN RESULT

THEOREM. *Let $V_1 \subset V_2$ be alternation systems of $C[a, b]$, $V_1, V_2 \subset C^1[a, b]$, $v_1 \in V_1$, and $v_2 \in V_2$. In order that there exists an element f in $C^1[a, b]$ such that $v_i \in P_{V_i}(f)$, $i = 1, 2$; it is necessary and sufficient that $v_1 - v_2$ is either zero or changes sign at atleast $d(v_1)$ points of $[a, b]$.*

For the proof we require the following lemmas.

LEMMA 1. Let M_1 be a compact Hausdorff space in \mathbb{R}^n and M_3 an open set containing M_1 . Then there exists a function F defined on M_3 with compact support $M_2 \supset M_1$ with the following properties:

- (a) $F \in C^\infty[M_3]$,
- (b) $F(t) = 1$ for $t \in M_1$,
- (c) $0 \leq F(t) \leq 1$ for $t \in M_2 \setminus M_1$.

For the proof of the above lemma see de Rham [4, p. 4].

LEMMA 2. Let U_1, U_{-1} be piecewise continuously differentiable functions in $C[a, b]$ and let

$$M_1 := \{t_1, t_2, \dots, t_k\},$$

$$M_{-1} := \{\tau_1, \tau_2, \dots, \tau_j\}$$

be disjoint subsets of the open interval (a, b) , with the following properties:

- (i) $U_{-1}(t) < 0 < U_1(t), \forall t \in [a, b]$.
- (ii) U_i is continuously differentiable at the points of $M_i, i = 1, -1$.

Then there exists an $f \in C^1[a, b]$ such that

- (a) $f(t_\mu) = U_1(t_\mu), \mu = 1, 2, \dots, k$,
- (b) $f(\tau_\nu) = U_{-1}(\tau_\nu), \nu = 1, 2, \dots, j$,
- (c) $U_{-1}(t) \leq f(t) \leq U_1(t), \forall t \in [a, b]$.

Proof. Set for $\mu = 1, 2, \dots, k$ and λ_1 , a positive real number,

$$G_\mu^{-1}(t, \lambda_1) := U_1(t) - \lambda_1(t - t_\mu)^2,$$

and for $\nu = 1, 2, \dots, j$ and λ_{-1} , a positive real number,

$$G_\nu^{-1}(t, \lambda_{-1}) := U_{-1}(t) + \lambda_{-1}(t - \tau_\nu)^2.$$

Then we have

$$G_\mu^{-1}(t_\mu, \lambda_1) = U_1(t_\mu), \quad \mu = 1, 2, \dots, k, \quad (1)$$

$$G_\nu^{-1}(\tau_\nu, \lambda_{-1}) = U_{-1}(\tau_\nu), \quad \nu = 1, 2, \dots, j. \quad (2)$$

Further we have that G_μ^{-1} and G_ν^{-1} are continuously differentiable at t_μ and τ_ν respectively. By choosing λ_1 and λ_{-1} small enough, we have that in an open interval (c_μ, d_μ) of t_μ and an open interval (c_ν^{-1}, d_ν^{-1}) of τ_ν ,

$$0 < G_\mu^{-1}(t, \lambda_1) \leq U_1(t), \quad \forall t \in (c_\mu, d_\mu), \quad (3)$$

$$0 > G_\nu^{-1}(t, \lambda_{-1}) \geq U_{-1}(t), \quad \forall t \in (c_\nu^{-1}, d_\nu^{-1}), \quad (4)$$

and G_μ^{-1} restricted to (c_μ, d_μ) and G_ν^{-1} restricted to (c_ν^{-1}, d_ν^{-1}) are, respectively, in $C^1(c_\mu, d_\mu)$ and $C^1(c_\nu^{-1}, d_\nu^{-1})$. We can further assume, by taking these intervals sufficiently small, that these intervals are pairwise disjoint. Now consider the interval (c_μ, d_μ) , $\mu = 1, 2, \dots, k$. Since $t_\mu \in (c_\mu, d_\mu)$, there exists a compact interval I_μ containing t_μ in its interior. Apply Lemma 1 with $M_1 = I_\mu$ and $M_3 = (c_\mu, d_\mu)$. Thus there exists a function F_μ and a compact set $\tilde{I}_\mu \supset I_\mu$ such that

- (ā) $F_\mu \in C^\infty(c_\mu, d_\mu)$,
- (b̄) $F_\mu(t) = 1$ for $t \in I_\mu$,
- (c̄) $0 \leq F_\mu(t) \leq 1$ for $t \in \tilde{I}_\mu \setminus I_\mu$.

We further define $F_\mu(t) = 0$ for $t \in [a, b] \setminus (c_\mu, d_\mu)$. Then it is clear that F_μ is in $C^\infty[a, b]$. Similarly we define for each ν , $\nu = 1, 2, \dots, j$; $F_\nu^{-1} \in C^\infty[a, b]$. We now set,

$$f(t) := \sum_{\mu=1}^k F_\mu(t) G_\mu^{-1}(t, \lambda_1) + \sum_{\nu=1}^j F_\nu^{-1}(t) G_\nu^{-1}(t, \lambda_{-1})$$

for all t in $[a, b]$. We claim that $f \in C^1[a, b]$ and satisfies (a), (b), and (c) of the lemma. Notice that f is well defined on $[a, b]$. Since $F_\mu(t) G_\mu^{-1}(t, \lambda_1) \in C^1[a, b]$ and $F_\nu^{-1}(t) G_\nu^{-1}(t, \lambda_{-1}) \in C^1[a, b]$, we have that $f \in C^1[a, b]$. If $t = t_\mu$, then from the definition of f it follows that

$$f(t_\mu) = F_\mu(t_\mu) G_\mu^{-1}(t_\mu, \lambda_1)$$

since for all the indices $i \neq \mu$, $F_i(t) = 0$ on (c_μ, d_μ) and for all indices ν , $\nu = 1, 2, \dots, j$; $F_\nu^{-1}(t) = 0$ on (c_μ, d_μ) . Since $F_\mu(t_\mu) = 1$,

$$f(t_\mu) = G_\mu^{-1}(t_\mu, \lambda_1) = U_1(t_\mu)$$

This proves (a). Similarly, when $t = \tau_\nu$, we have $f(\tau_\nu) = U_{-1}(\tau_\nu)$. Finally, if $t \in (c_\mu, d_\mu)$, then, $0 \leq F_\mu(t) \leq 1$ and from (3), we have

$$U_{-1}(t) < 0 \leq f(t) = F_\mu(t) G_\mu^{-1}(t, \lambda_1) \leq U_1(t).$$

Similarly, if $t \in (c_\nu^{-1}, d_\nu^{-1})$, we have $0 \leq F_\nu^{-1}(t) \leq 1$, and from (4) it follows that

$$U_1(t) > 0 \geq f(t) = F_\nu^{-1}(t) G_\nu^{-1}(t, \lambda_{-1}) \geq U_{-1}(t).$$

If $t \notin \{\bigcup_{\mu=1}^k (c_\mu, d_\mu)\} \cup \{\bigcup_{\nu=1}^j (c_\nu^{-1}, d_\nu^{-1})\}$, then clearly we have that

$$U_{-1}(t) < f(t) < U_1(t).$$

Combining these we have

$$U_{-1}(t) \leq f(t) \leq U_1(t) \quad \text{for all } t \in [a, b].$$

This completes the proof of Lemma 2.

Proof of the theorem. The proof is exactly the same as it is for the case when $V_1 \subset V_2 \subset C^2[a, b]$ (see Brosowski [2]). For the sake of completeness we give here the proof of the sufficiency part. The case $v_1 = v_2$ follows from Brosowski [1, Theorem 2.15, p. 62]. Let $v_1 \neq v_2$. Let $a_1, a_2, \dots, a_{d(v_1)}$ be the zeros of $v_1 - v_2$ taken in their natural order. Put $a_0 = a$ and $a_{d(v_1)} + 1 = b$. Then we have

$$a = a_0 < a_1 < a_2 < \dots < a_{d(v_1)} < a_{d(v_1)} + 1 = b.$$

Then there exists an $\eta \in \{-1, 1\}$ and for $i = 0, 1, 2, \dots, d(v_1)$ points $t_{1,i}$ in (a_i, a_{i+1}) such that

$$\eta(-1)^i(v_2(t_{1,i}) - v_1(t_{1,i})) \geq \beta > 0$$

for $i = 0, 1, \dots, d(v_1)$. Since $v_2 - v_1$ has at least one zero in (a, b) , we can choose $d(v_2) + 1$ points $t_{2,0}, t_{2,1}, \dots, t_{2,d(v_2)}$ such that

$$a \leq t_{2,0} < t_{2,1} < \dots < t_{2,d(v_2)} \leq b$$

and

$$|v_2(t_{2,i}) - v_1(t_{2,i})| < \beta/2$$

for $i = 0, 1, \dots, d(v_2)$. With a suitable real number $A > 0$, we set

$$U_1(t) := \min\{(v_1(t) + A + \beta), (v_2(t) + A + \beta/2)\} \quad (\text{x})$$

$$U_{-1}(t) := \max\{(v_1(t) - A - \beta), (v_2(t) - A - \beta/2)\} \quad (\text{xx})$$

Then we have, if A is sufficiently large, that

$$U_{-1}(t) < 0 < U_1(t) \quad \text{for all } t \in [a, b].$$

Notice that at a point $t_{1,i}$ with $\eta(-1)^i = +1$, we have

$$v_1(t_{1,i}) + A + \beta < v_2(t_{1,i}) + A + \beta/2.$$

Hence in a neighborhood of $t_{1,i}$ also we have the same inequality. Similarly,

at a point $t_{1,s}$ with $\eta(-1)^s = -1$, we have that if $t \in$ a small neighborhood of $t_{1,s}$,

$$v_1(t) - A - \beta > v_2(t) - A - \beta/2.$$

This shows that U_1 is continuously differentiable at the points $t_{1,i}$ with $\eta(-1)^i = 1$ and U_{-1} is continuously differentiable at the points $t_{1,s}$ with $\eta(-1)^s = -1$. Similarly we can show that at the points $t_{2,i}$, with i even, U_1 is continuously differentiable and U_{-1} is so at the points $t_{2,s}$ with s odd. Thus if we set

$$\begin{aligned} M_1 &:= \{t_{1,i} \text{ with } \eta(-1)^i = 1\} \cup \{t_{2,i} \text{ with } i \text{ even}\} \\ M_{-1} &:= \{t_{1,s} \text{ with } \eta(-1)^s = -1\} \cup \{t_{2,s} \text{ with } s \text{ odd}\}, \end{aligned}$$

we have that U_i is continuously differentiable on M_i , $i = 1, -1$. By Lemma 2 we have that there exists an $f \in C^1[a, b]$ such that

- (i) $U_{-1}(t) \leq f(t) \leq U_1(t)$ for all $t \in [a, b]$,
- (ii) $f(t) = U_1(t)$ on M_1 ,
- (iii) $f(t) = U_{-1}(t)$ on M_{-1} .

Now from (x) and (xx) it follows that

$$\begin{aligned} -(A + \beta) &\leq f(t) - v_1(t) \leq A + \beta \\ -(A + \beta/2) &\leq f(t) - v_2(t) \leq A + \beta/2 \end{aligned}$$

for all $t \in [a, b]$. If $t = t_{1,i}$ with $\eta(-1)^i = 1$ then we have

$$f(t_{1,i}) = U_1(t_{1,i}) = v_1(t_{1,i}) + A + \beta$$

and so we have

$$f(t_{1,i}) - v_1(t_{1,i}) = A + \beta$$

while if $t = t_{1,s}$ with $\eta(-1)^s = -1$ we have

$$f(t_{1,s}) - v_1(t_{1,s}) = -A - \beta.$$

Similarly if $t = t_{2,i}$ with i even, we have

$$f(t_{2,i}) - v_2(t_{2,i}) = A + \beta/2$$

and if $t = t_{2,s}$ with s odd we have

$$f(t_{2,s}) - v_2(t_{2,s}) = -A - \beta/2.$$

Since V_1 and V_2 are alternation systems we conclude that

$$v_i \in P_{V_i}(f), \quad i = 1, 2.$$

This completes the proof of the theorem.

ACKNOWLEDGMENTS

The author wishes to record his gratitude to Professor Dr. B. Brosowski for suggesting this problem and for useful discussions and to the Humboldt Foundation for financial support. His thanks are also due to the referee for useful comments.

REFERENCES

1. B. BROSWOSKI, "Nicht-Lineare Tschebyscheff-Approximation," Hochschulschriften, Bd. 808–808a, Bibliographisches Institut, Mannheim, 1968.
2. B. BROSWOSKI, On simultaneous best approximation, to appear.
3. B. BROSWOSKI AND M. R. SUBRAHMANYA, On the existence of functions with prescribed best approximations, *J. Approximation Theory* **15** (1975), 143–155.
4. G. DE RHAM, "Variétés Différentiables," Hermann, Paris, (1955).
5. F. DEUTSCH, P. D. MORRIS AND I. SINGER, On a problem of T. J. Rivlin in approximation theory, *J. Approximation Theory* **2** (1969), 342–352.
6. H. G. HEGERING, Über das Rivlin-Problem der Simultanen Inversen Tschebyscheff-Approximation, Thesis, Universität München, 1971.
7. T. J. RIVLIN, New and Unsolved Problems, No. 14: Best Algebraic Approximation, Abstrakte Räume und Approximation, Vol. 10, P. 421, Birkhäuser-Verlag, Basel/Stuttgart, 1969.
8. D. A. SPRECHER, Simultaneous best approximations with two polynomials, *J. Approximation Theory*, **2** (1969), 384–388.
9. D. A. SPRECHER, On simultaneous Chebyshev approximations, *J. Approximation Theory* **4** (1971), 137–146.
10. M. R. SUBRAHMANYA, A note on a problem of Rivlin, *J. Approximation Theory* **6** (1972), 359–361.
11. M. R. SUBRAHMANYA, A complete solution to a problem of T. J. Rivlin, to appear.